

(3)

RR-82-33-ONR

SAMPLING VARIANCES AND COVARIANCES
OF PARAMETER ESTIMATES IN
ITEM RESPONSE THEORY

Frederic M. Lord and Marilyn S. Wingersky

This research was sponsored in part by the Personnel and Training Research Programs Psychological Sciences Division Office of Naval Research, under Contract No. N00014-80-C-0402

Contract Authority Identification Number NR No. 150-453

Frederic M. Lord, Principal Investigator



Educational Testing Service Princeton, New Jersey

August 1982



Reproduction in whole or in part is permitted for any purpose of the United States Government.

Approved for public release; distribution unlimited.

SAMPLING VARIANCES AND COVARIANCES

OF PARAMETER ESTIMATES IN

ITEM RESPONSE THEORY

Frederic M. Lord and Marilyn S. Wingersky

This research was sponsored in part by the Personnel and Training Research Programs Psychological Sciences Division Office of Naval Research, under Contract No. N00014-80-C-0402

Contract Authority Identification Number NR No. 150-453

Frederic M. Lord, Principal Investigator


Educational Testing Service Princeton, New Jersey

August 1982



Reproduction in whole or in part is permitted for any purpose of the United States Government.

Approved for public release; distribution unlimited.



UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION	READ INSTRUCTIONS BEFORE COMPLETING FORM	
1 REPORT NUMBER	2. 000   4002331011 110.	
	AD-A119609	
4 TITLE (and Subtitle)		5 TYPE OF REPORT & PERIOD COVERED
Sampling Variances and Covariances of Parameter Estimates in Item Response Theory		Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7 AuThoris,		RR-82-33-ONR  B CONTRACT OR GRANT NUMBER(a)
Frederic M. Lord and Marilyn S. Win	ngersky	N00014-80-C-0402
PERFORMING ORGANIZATION NAME AND ADDRESS Educational Testing Service Princeton, NJ 08541		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  NR 150-453
Personnel and Training Research Proffice of Naval Research (Code 458	•	12 REPORT DATE August 1982
Arlington, VA 22217	5)	13 NUMBER OF PAGES
4 MONITURING AGENCY NAME & ADDRESS(II dilferent	from Controlling Office)	15. SECURITY CLASS rot this report:
		Unclassified  15. DECLASSIFICATION DOWNGRADING SCHEDULE
Approved for public release; disti		
18 SUPPLEMENTARY NOTES		
19 KEY WORDS (Continue on reverse side if necessary and	identify by block number)	
Sampling Covariances Standard Errors Item Response Theory Maximum Likelihood Estimators	Item Parameters Matrix Inversion	•
This paper develops a possible pling variance-covariance matrix item response theory when both ite For a set of artificial data, results with the variance-covariance case where the abilities are known the results are consistent with the asymptotically correct except for	e method for composition of joint maximum of joint maximum em parameters and alts are compared matrices found in, or where the ne conjecture that	likelihood estimates in d abilities are unknown. d with empirical values; by the usual formulas for the item parameters are known. at the new method is
DD (1975) 1473 FORTH NOF FINGUES IS OBSOLE NO STORES TO SERVE T	UNCLAS SECTION OF AS	SIFIED SSIFICATION OF THIS PAGE (When Date Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)	
	}
	Ì
	[
	}
	į
	1
	ł
	Ì
	1
	1
	1
	1
	1
	1
	1
	1
	I
	j
	1
	ļ
	İ
	İ
	İ
	1
	[
	- 1
	1
	- 1

# Sampling Variances and Covariances of Parameter Estimates in Item Response Theory

#### Abstract

This paper develops a possible method for computing the asymptotic sampling variance-covariance matrix of joint maximum likelihood estimates in item response theory when both item parameters and abilities are unknown. For a set of artificial data, results are compared with empirical values; also with the variance-covariance matrices found by the usual formulas for the case where the abilities are known, or where the item parameters are known. The results are consistent with the conjecture that the new method is asymptotically correct except for errors due to grouping.

## Sampling Variances and Covariances of Parameter Estimates in Item Response Theory\*

In item response theory (IRT), the observations come in the form of an n-by- N matrix, with one row for each item and one column for each examinee. The joint frequency distribution of the observations depends on a vector of N 'ability' parameters—one for each person—and on a matrix of item parameters. Here, we will consider only the three-parameter logistic model for dichotomously scored items, so there will be three item parameters ( a , b , and c ) for each of n items. A method will be developed for computing the asymptotic sampling variance—covariance matrix when both abilities and item parameters are unknown. Until this is done we do not know the standard errors of the parameter estimates, which handicaps development of a goodness—of fit test and other statistics required in applications of IRT.

If the item (ability) parameters are known, the estimated ability (item) parameters have independent sampling distributions. It can be shown (see Bradley & Gart, 1962) that the maximum likelihood estimates of the ability (item) parameters are consistent. Hence the asymptotic sampling variance for an estimated ability parameter is given by the usual formula

$$Var(\{\frac{1}{r}, a, b, c\}) = \left[S(\frac{1}{r}, c)^2\right]^{-1} , \qquad (1..)$$

where  $\frac{1}{r}$  is the estimated ability parameter,  $\ell$  is the log of the likelihood, and a, b, and c are the known vectors of item parameters.

<sup>\*</sup>This work was supported in part by contract N00014-80-C-0402, project designation NR 150-453 between the Office of Naval Research and Educational Testing Service. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Similarly the asymptotic sampling variance-covariance matrix of the estimated item parameters for an item is given by

$$\|\operatorname{Cov}(\hat{\tau}_{\mathbf{v}},\hat{\tau}_{\mathbf{w}},\hat{\tau})\| = \|\mathcal{E}(\frac{\partial \mathbf{f}}{\partial \tau_{\mathbf{v}}} \frac{\partial \mathbf{f}}{\partial \tau_{\mathbf{w}}})\|^{-1} \qquad (v,w=1,2,3)$$

where  $\frac{1}{V}$  is a vector consisting of the estimated a , b , and c for a single item and — is the known vector of abilities. The right-hand side is the inverse of a 3-bv-3 matrix.

When neither item nor ability parameters are known, all parameters are often estimated simultaneously by maximum likelihood. In the (Rasch) case where there is only one parameter per item, Haberman (1977) has shown that all parameter estimates will converge to their true values (will be consistent) when the number of examinees and the number of test items become large simultaneously. Empirical results suggest that consistency probably also holds when all parameters are estimated simultaneously under the three-parameter model. If so, it is reasonable that the asymptotic sampling variance-covariance matrix of all estimated parameters will be given by the usual formula

$$\|\operatorname{Cov}(\tau_{\mathbf{p}}, \tau_{\mathbf{q}})\| = \|\mathfrak{s}(\frac{2\mathbf{\ell}}{\tau_{\mathbf{p}}}, \frac{2\mathbf{\ell}}{2\tau_{\mathbf{q}}})\|^{-1} \qquad (\mathbf{p}, \mathbf{q} = 1, 2, ..., \mathbf{M})$$

where M = 3n + N - 2 and  $E = \frac{1}{p} \cdot E \cdot a_1, b_1, c_1, a_2, b_2, c_2, \dots, a_n, b_n, c_n;$   $\frac{1}{2}, \dots, \frac{1}{N-2}, \dots$ 

Since standard errors are urgently needed in practical work where all parameters are estimated simultaneously by maximum likelihood, this report compares numerical values provided by (2) with values provided by (1) and with empirically observed sampling fluctuations. The comparisons to be presented suggest that (2) provides useful values for the desired standard errors.

There are several special problems that arise in the evaluation and practical utilization of (2), problems that do not arise in the situation where (1) is appropriate:

- Until an origin and scale are specified, the parameters are not identifiable.
- 2. The mathematical formulation is complicated by the choice of origin and scale.
- 3. The usual choice of origin and scale when estimating IRT parameters is inconvenient for mathematical purposes.
- 4. The numerical values of the sampling variances are very much affected by the choice of origin and scale.
- 5. Equation (2) requires the inversion of a matrix of order N+3n-2 where N may be several thousand.

These problems will be considered in subsequent sections.

## 1. Parameterization

The appropriate likelihood function is (Lord, 1980)

$$L(\mathbf{a}, \mathbf{b}, \mathbf{c}; \mathbf{a}|\mathbf{U}) = \frac{\mathbf{n}}{\mathbf{n}} \sum_{i=1}^{N} \mathbf{p}_{ia}^{\mathbf{u}} \mathbf{a}_{0}^{\mathbf{1}-\mathbf{u}} \mathbf{i} \mathbf{a}$$
(3)

where \_\_is the vector of the N\_ability parameters; a\_, b\_, and c\_are each a vector of \_n\_item parameters,  $\Gamma \equiv u_{ia}$  \_is the matrix of item responses  $u_{ia}$  (= 0 or 1); finally  $Q_{ia} \equiv 1 - P_{ia}$  and  $P_{ia}$  is the item response function, the probability of a correct answer by examinee a\_to item\_i . Each given  $P_{ia}$  is a function of \_a\_a and of \_a\_i , b\_i , and \_c\_i , but not of any other parameters. In numerical work here,  $P_{ia}$  will be taken to be the three-parameter logistic function

$$P_{ia} = c_{i} + \frac{1 - c_{i}}{1 + \exp[-1.7a_{i}(\theta_{a} - b_{i})]} . \tag{4}$$

For mathematical purposes, however, it is only necessary to state that  $P_{\underline{i}\underline{a}} \quad \text{is an increasing function of} \quad \frac{1}{a} \ .$ 

If we add some constant to all  $\frac{1}{a}$  and subtract the same constant from all  $b_i$ , all  $P_{ia}$  will be unchanged. This means that the origin used for measuring ability is entirely arbitrary. If we multiply each  $\frac{1}{a}$  and each  $\frac{1}{b}$  by some constant and divide each  $\frac{1}{a}$  by the same constant, again all  $P_{ia}$  will be unchanged. This means that the unit used to measure ability is entirely arbitrary. Since we can change the origin and unit of the  $\frac{1}{a}$  without changing (3), it follows that  $\frac{1}{a}$ ,  $\frac{1}{a}$ , and  $\frac{1}{a}$  are not identifiable and cannot be estimated from (3) without further specification.

To conform to a commonly used procedure, we could choose the origin and scale so that for some specified group of examinees the mean of the  $\frac{1}{a}$  is zero and the variance is one. This is not convenient mathematically, however. Instead, two other methods of

specifying the origin and scale will be used, even though this will complicate matters later on when the results are applied in practice. In the first method, without loss of generality, arbitrary numerical values will be assigned to  $\theta_{N-1}$  and to  $\theta_{N}$ .

The  $M \equiv N + 3n - 2$  likelihood equations are

$$0 = \sum_{i=1}^{n} \sum_{a=1}^{N} (u_{ia} - P_{ia}) \frac{P_{ia}^{ia}}{P_{ia}^{0}} \qquad (p = 1, 2, ..., M)$$
 (5)

where  $P_p^{ia} \equiv \Im P_{ia}/\Im \tau_p$ .

## 2. Fisher Information Matrix

The Fisher information matrix on the right of (2) now has as  ${\bf a}$  typical element

$$I_{pq} = \mathcal{E}(\frac{\partial \mathcal{L}}{\partial \tau_{p}} \frac{\partial \mathcal{L}}{\partial \tau_{q}}) = \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{a=1}^{N} \sum_{b=1}^{N} \frac{P_{i}^{1a}P_{j}^{1b}}{P_{ia}Q_{ia}P_{jb}Q_{jb}} Cov(u_{ia}, u_{jb})$$

$$(p,q=1,2,...,M)$$

Because of local independence and random sampling of examinees,

$$Cov(u_{ia}, u_{jb}) = \delta_{ij} \delta_{ab}^{P}_{ia} Q_{ia}$$

where  $\delta_{st} = 1$  if s = t ,  $\delta_{st} = 0$  otherwise. Thus the typical element is

$$I_{pq} = \frac{n}{i=1} \frac{N}{a=1} \frac{p}{p} \frac{q}{Q_{1a}} \quad (p,q=1,2,...,M) \quad . \tag{6}$$

Note that  $p^{ia}$  is zero unless either p and a refer to the same person, or p and i refer to the same item. Thus

$$\|\mathbf{I}_{pq}\| = \begin{bmatrix} \mathbf{S}_{1} & 0 & \dots & 0 & | & \mathbf{f}_{11} & \mathbf{f}_{12} & \dots & \mathbf{f}_{1N}, \\ 0 & \mathbf{S}_{2} & \dots & 0 & | & \mathbf{f}_{21} & \mathbf{f}_{22} & \dots & \mathbf{f}_{2N}, \\ \vdots & \vdots & & \vdots & & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & \mathbf{S}_{n} & | & \mathbf{f}_{n1} & \mathbf{f}_{n2} & \dots & \mathbf{f}_{nN}, \\ \vdots & \vdots & & \vdots & & \vdots & & \vdots & \ddots & \vdots \\ \mathbf{f}_{11}^{\prime} & \mathbf{f}_{21}^{\prime} & \dots & \mathbf{f}_{n2}^{\prime} & | & 0 & \mathbf{t}_{2} & \dots & 0 \\ \vdots & \vdots & & \vdots & & \vdots & & \vdots & \vdots \\ \mathbf{f}_{N}^{\prime}, \mathbf{f}_{N}^{\prime}, 2 & \dots & \mathbf{f}_{N}^{\prime}, n & | & 0 & 0 & \dots & \mathbf{t}_{N}, \end{bmatrix}$$

$$(7)$$

where N'  $\equiv$  N ~ 2 , S<sub>i</sub> is the 3-by-3 Fisher information matrix for  $a_i$ ,  $b_i$ , and  $c_i$ ,  $t_a$  is the Fisher information for examinee a, and  $f_{ia}$  is the 3-by-1 joint Fisher information vector for item i and examinee a:

$$f_{ia} = \frac{\frac{\partial P_{ia}}{\partial a}}{\frac{\partial P_{ia}}{\partial a}} \begin{bmatrix} \frac{\partial P_{ia}}{\partial a} \\ \frac{\partial P_{ia}}{\partial b} \\ \frac{\partial P_{ia}}{\partial c} \end{bmatrix}.$$

## 3. Matrix Inversion

The following general formula for inverting a partitioned matrix may be applied to (7)

$$\begin{bmatrix} S & F \\ F' & T \end{bmatrix}^{-1} = \begin{bmatrix} S^{-1} + S^{-1}FZ^{-1}F'S^{-1} & -S^{-1}FZ^{-1} \\ -Z^{-1}F'S^{-1} & Z^{-1} \end{bmatrix}$$
(8)

where

$$7 \equiv T - F'S^{-1}F \qquad .$$

The matrix S is easily inverted since it is a diagonal supermatrix:

$$S^{-1} = \| S_i^{-1} \|^{c}$$
.

The notation on the right denotes a diagonal matrix with diagonal elements  $\mathbf{S}_i^{-1}$ . These last are easily computed since each  $\mathbf{S}_i$  is only 3 by 3.

All the matrix operations indicated on the right side of (8) can be carried out on the computer without difficulty, with one exception: the inversion of Z , which is N' by N'. The approximation used here to invert Z relies on grouping the  $\theta_a$  into 16 class intervals of width 0.5, covering the range  $-5 \leq \theta_a \leq 3$ . Each  $\theta_a$  in a given class interval is replaced by the midpoint of the interval.

Now T will be a diagonal supermatrix  $T \equiv \begin{bmatrix} T \\ 0 \end{bmatrix}$ , where  $T_g \equiv t_g I$  is a scalar matrix with dimensions  $N_g$  by  $N_g$ , and  $N_g$  is the number of people in class interval g. Also, F will be a row vector of 16 matrices, the columns of any one matrix being all identical:

$$F \equiv \{f_1 1'_1, f_2 1'_2, \dots, f_{16} 1'_{16}\} \quad , \tag{10}$$

where  $f(g) \equiv \{f(g)\}$  for any examinee a in class interval g and f(g) is a unit vector whose length is  $N_g$  .

The product  $F'S^{-1}F$  can now be written as a 16-by-16 supermatrix:

$$F'S^{-1}F = \{[1_{g-g}f'S^{-1}f_{h-h}]\}$$

Denote the scalar  $f'_{g}S^{-1}f_{h}$  by  $w_{gh}$  . We now have

$$z \equiv T - \lim_{gh^{(i)}}, \qquad (11)$$

$$M_{gh} \equiv w_{gh} \frac{1}{8} \frac{1}{h} . \tag{12}$$

For computation purposes, Z still has N' rows and columns, not just 16. For the usual sample size, it is still not feasible to invert Z with a standard inversion program.

Consider the problem of inverting  $z_{11}$ , the  $N_1$ -by- $N_1$  upper left corner of z. By (11), (12), and a standard formula,

$$z_{11}^{-1} = \left[ T_1 - w_{11}^{-1} 1_{11}^{-1} \right]^{-1} = T_1^{-1} + \frac{w_{11}^{-1} 1_{11}^{-1} 1_{11}^{-1}}{1 - w_{11}^{-1} 1_{11}^{-1} T_1^{-1}} . \tag{13}$$

Since  $T_1 \equiv t_1 I$ , where  $t_1$  is scalar, this becomes

$$Z_{11}^{-1} = \frac{1}{t_1} + \frac{w_{11}^{1} \frac{1}{1} \frac{1}{1}}{t_1^{2} - t_1 w_{11}^{N_1}}.$$

Next, the upper left 2-by-2 supermatrix in % can be inverted as in (8), using the standard formula for the inversion of a partitioned matrix:

$$\begin{bmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \end{bmatrix}^{-1} = \begin{bmatrix} z_{11}^{-1} + z_{11}^{-1} z_{12} H^{-1} z_{21} z_{11}^{-1} & -z_{11}^{-1} z_{12} H^{-1} \\ -H^{-1} z_{21} z_{11}^{-1} & H^{-1} \end{bmatrix}$$
(14)

where  $H = Z_{22} - Z_{21}Z_{11}^{-1}Z_{12}$ . It can be seen that H has the same general form as  $Z_{11}$  and can thus be inverted as in (13); so (14) can readily be calculated.

Next, substitute (14) for  $\mathbb{Z}_{11}^{-1}$  in the foregoing procedure, and repeat this procedure, in such a way as to invert the upper

left 3-by-3 supermatrix in  $\mathbb{Z}$ . A total of fifteen repetitions enable us to invert the 16-by-16 supermatrix  $\mathbb{Z}$ . Equation (8) is now used for one final inversion, the result being the desired variance-covariance matrix of all  $\mathbb{N} + 3n - 2$  parameters.

The 16-by-16 variance-covariance supermatrix for the consists of 256 blocks. The elements are all the same within a block except for diagonal blocks, each of which has a variance (instead of a covariance) repeated along its diagonal. Any two examinees in the same class interval will have identical Var and identical sampling covariances with any other given parameter estimate.

#### 4. Reparameterization

In Section 1, in order to have identifiable parameters, an origin and scale was chosen so that  $\frac{1}{N-1}$  and  $\frac{1}{N}$  had arbitrary preassigned values. Any other choice of origin and scale would result in a linear transformation of parameters. The likelihood function would remain unchanged for every pattern of item responses.

The choice of unit (but not the choice of origin) has one completely obvious effect on the sampling errors of parameter estimates. If the unit is changed, the standard errors for the 's and 's will be multiplied by the ratio of the new scale unit to the old scale unit. The standard errors for the a 's will be divided by this ratio. A second important effect is easily overlooked: the standard error

of the maximum likelihood estimator depends not only on the choice of scale, but also on how the (origin and) scale is specified.

Suppose that the <u>true</u> numerical values of all  $\theta_a$  (a = 1,...,N) are specified on some arbitrary scale. Suppose next that our test is too difficult for examinee N. This means that the likelihood function is rather insensitive to variations in  $\theta_N$ . If we could repeat our testing with several parallel test forms, we would find a wide range of estimates of  $\theta_N$ . In such a situation, the difference between true  $\theta_{N-1}$  and  $\theta_N$  clearly cannot be estimated well from the examinee responses. If we define the scale by treating  $\theta_N$  and  $\theta_{N-1}$  as known, our estimates of every  $\theta_A$  may fluctuate grossly, simply because the scale unit  $\theta_N - \theta_{N-1}$  is not well determined by the data.

Suppose next that we relabel all examinees so that examinees N-1 and N are not the same examinees as before. The ability scale has not been changed from the preceding paragraph; it is the procedure for defining the scale that has been changed. The true  $^{\circ}$  for each examinee is still the same as before. Suppose the new examinees N-1 and N are both at ability levels where our test measures accurately. If, further, the true  $\frac{1}{N-1}$  and  $\frac{1}{N}$  are substantially different from each other, the difficulty of the previous paragraph disappears: Throughout the ability range where the test is designed to measure accurately, the standard errors of all  $\frac{1}{N}$  may be reasonably small.

For example, suppose on some scale  $\theta_1 = -3$ ,  $\theta_2 = -2$ ,  $\theta_3 = -1$ ,  $\theta_4 = 0$ ,  $\theta_5 = 1$ ,  $\theta_6 = 2$ ,  $\theta_7 = 3$ . We can specify this same scale in terms of any two of these  $\theta$  's. The standard errors that we obtain will depend in an overwhelming way not just on the ability scale, but on how we specify it. We cannot rectify the standard errors by some simple procedure, such as multiplying each by a constant.

For this reason, our procedure for specifying the ability scale should depend only on parameters or functions of parameters that are accurately determined by the data. A robust mean of the  $\theta_a$  might seem attractive; however, any function of the  $\theta_a$  is counterindicated by the fact that sometimes  $\hat{\theta}_a = \pm \infty$ .

The procedure used here is to choose a set of m discriminating, moderately easy items and a set of r discriminating, moderately hard items. We will hereafter define the origin and unit for our new parameters, to be denoted by capital letters, so that the mean of the (true) B -parameters for the easy items is zero, and the mean for the hard items is one.

Our new parameters are related to our old parameters (from Section 2 or from Section 5) by linear transformations:

$$A_{i} \equiv ka_{i}$$
,  $B_{i} \equiv K + b_{i}/k$ ,  $C_{i} \equiv C_{i}$ ,  $O_{a} \equiv K + O_{a}/k$ , (15)
$$(a = 1, 2, ..., N; i = 1, 2, ..., n)$$

where k and K are transformation constants to be determined. Since

$$\bar{\mathbf{B}}_0 \equiv \frac{1}{m} \stackrel{m}{\Sigma} \mathbf{B}_i = 0 , \quad \bar{\mathbf{B}}_1 \equiv \frac{1}{r} \stackrel{r}{\Sigma} \mathbf{B}_i = 1 , \qquad (16)$$

the values of  $\,k\,$  and  $\,K\,$  are found by substituting (15) into (16) and solving for  $\,k\,$  and  $\,K\,$ :

$$\mathbf{k} = \overline{\mathbf{b}}_1 - \overline{\mathbf{b}}_0 , \qquad \mathbf{K} = -\frac{\overline{\mathbf{b}}_0}{\mathbf{k}} , \qquad (17)$$

where  $\bar{b}_0$  and  $\bar{b}_1$  are means for m and r items, respectively.

To find the variance-covariance matrix for estimates of the uppercase parameters, rewrite (15) as

$$\Theta_{a} = (\theta_{a} - \bar{b}_{0})/k$$
,  $A_{i} = ka_{i}$ ,  $B_{i} = (b_{i} - \bar{b}_{0})/k$ , (18)  
 $C_{i} = c_{i}$ .

Because of the special properties of maximum likelihood estimators, equations (18) still hold when estimators are substituted for parameters. Thus the sampling variances and covariances for estimates of the new parameters can be computed from the sampling variances and covariances already obtained at the end of Section 3. Formulas for doing this can be written down from (18) by using the 'delta' method (Kendall & Stuart, 1969, Chapter 10). For example,

$$\operatorname{Cov}(\hat{A}_{i}, \hat{b}_{0}) = \operatorname{Cov}(\hat{a}_{i}, \hat{b}_{0}) - \operatorname{Cov}(\hat{a}_{i}, \hat{b}_{0}) - \frac{\hat{a}_{i} - \hat{b}_{0}}{k} \operatorname{Cov}(\hat{a}_{i}, \hat{k}) + \frac{a_{i}}{k} \operatorname{Cov}(\hat{b}_{0}, \hat{k}) - \frac{a_{i}}{k} \operatorname{Cov}(\hat{b}_{0}, \hat{k}) - \frac{a_{i}(\theta_{a} - \hat{b}_{0})}{k^{2}} \operatorname{Var} k ,$$

$$\operatorname{Cov}(\hat{b}_{0}, \hat{k}) = \operatorname{Cov}(\hat{b}_{1}, \hat{b}_{0}) - \operatorname{Var} \hat{b}_{0} ,$$

$$\operatorname{Cov}(\hat{b}_{1}, \hat{b}_{0}) = \frac{1}{mr} \sum_{i} \operatorname{Cov}(\hat{b}_{i}, \hat{b}_{i}) .$$

## 5. Parameter Estimation

The maximum likelihood estimators (MLE) satisfy the likelihood equations (5). In (5), there is one equation for each parameter omitting  $\theta_{N-1}$  and  $\theta_N$ . If all N + 3n  $\equiv$  M + 2 MLE are linearly transformed, as for example in (15), the transformed parameters will still satisfy the likelihood equations.

Since the origin and scale for the new parameters is chosen to satisfy (16), then the appropriate k and K are obtained from (17) after replacing  $\bar{b}_0$  and  $\bar{b}_1$  by their MLE. The likelihood function (3) is unaffected by these linear transformations.

The computer program LOGIST identifies the parameters by still another choice of origin and scale:

- 1. a certain truncated mean of the  $\hat{\theta}_a$  ( a = 1,2,...,N ) is set equal to zero,
- 2. a certain truncated standard deviation of the  $\frac{1}{a}$  is set equal to one.

We will use the usual lower case symbols for parameters on this LOGIST scale. This should not cause confusion, since the lower-case parameters of Sections 1-3 will not be needed again.

If we start with LOGIST  $\hat{a}_i$ ,  $\hat{b}_i$ ,  $\hat{c}_i$ , and  $\hat{\theta}_a$  and determine k and K so that  $\hat{B}_0 = 0$  and  $\hat{B}_1 = 1$ , then the  $\hat{A}_i$ ,  $\hat{B}_i$ ,  $\hat{C}_i$  (i = 1, 2, ..., n), and the  $\hat{\theta}_a$  (a = 1, 2, ..., N), calculated by substituting estimated values into (15), will still satisfy the likelihood equations. The upper-case parameter estimates so obtained should have the sampling variance-covariance matrix found theoretically at the end of Section 4. Our remaining task is to compare an empirically determined variance-covariance matrix of MLE's with the corresponding theoretical matrix.

#### 6. Recapitulation

We have used, at different points, three different arbitrary scales for our parameters:

- 1.  $e_N$  and  $\theta_{N-1}$  are assigned arbitrarily.
- 2. The origin is set at  $\overline{B}_0$  , the unit is  $\overline{B}_1$  .

3. The origin is set at a truncated mean of the  $\frac{1}{a}$ , the unit is a truncated standard deviation of the

Scale 1 (denoted by lower-case symbols) is most convenient mathematically for the difficult task of inverting the M-by-M information matrix. Scale 1 is not useful for practical purposes, however, since its use grossly inflates all the sampling variances.

Scale 2 (denoted by upper-case symbols) seems the simplest choice in an attempt to keep the sampling error in the estimated origin and unit as small as possible. The sampling variances computed for scale 1 are transformed (see eq. 19) to values appropriate for scale 2. Although scale 2 is not the familiar one, the two item sets used to specify the scale can be chosen so that the numerical values of  $\hat{A}_i$ ,  $\hat{\beta}_i$ ,  $\hat{C}_i$  differ little from the familiar  $\hat{a}_i$ ,  $\hat{b}_i$ , and  $\hat{c}_i$  produced by LOGIST.

Scale 3 (hereafter denoted by lower-case symbols) is the scale used by LOGIST.

## 7. Empirical Estimation Procedures

As already stated, our theoretical results can be trusted only if they are shown to be in reasonable agreement with empirical results. For this purpose, artificial data  $\|\mathbf{u}_{ia}\|$  were created representing the administration of a 45-item test to a random sample of 1500

examinees. The 1500  $e_a$  were a spaced sample drawn from a distribution of abilities from a regular test administration. Six replicate matrices of  $\|\mathbf{u}_{ia}\|$  were independently generated, using the same item parameters and the same 1500  $e_a$ . The variation in responses across these matrices thus represents random fluctuations in  $\mathbf{u}_{ia}$  for fixed  $\mathbf{a}_i$ ,  $\mathbf{b}_i$ ,  $\mathbf{c}_i$  and  $e_a$ .

Further replication was also built in: items 16-30 and items 31-45 had the same item parameters as items 1-15. The true lower-case and upper-case item parameters are shown in Table 1 for items 1-15.

Six independent runs were made on LOGIST, one for each group of 1500 examinees. For each run separately,  $\hat{b}_0$  was calculated from items 4-9, 19-24, 34-39;  $\hat{b}_1$  was calculated from items 10-15, 25-30, 40-45. It is convenient for our ultimate interpretation of the standard errors to be obtained that the true  $\hat{b}_1 - \hat{b}_0 = .671 - (-.305) = .976$ . Since this is close to 1.0, the scale unit for the capitalized parameters is very close to the scale unit for the lower-case (LOGIST) parameters.

For each run separately, all lower-case parameter estimates were linearly transformed as in (15) to the upper-case scale, using estimated k and K values. For the data reported in subsequent sections, the true k = .976 and the true K = .312 . Since the six runs are independent, an unbiased empirical estimate of the sampling variance of any parameter estimate T is given by

Table 1
True (Upper Case) Item Parameters

Item					С
No.	<u>A</u>	a	<u>B</u>	<u>b</u>	or c
1	.96	.99	-1.75	-2.01	.17
2	.34	.35	-1.33	-1.61	.17
3	1.34	1.38	~.80	-1.09	.17
4	.76	.78	48	<b></b> 77	.17
5	.41	.42	38	67	.17
6	.90	. 92	04	34	.17
7	.90	. 92	.16	<b></b> 15	.17
8	1.04	1.06	.31	.00	.17
9	1.31	1.34	.42	.11	.13
10	1.46	1.50	.58	.26	.34
11	.85	.87	.79	.46	.17
12	.60	.62	.90	.57	.17
13	1.06	1.09	1.01	.68	.25
14	1.36	1.39	1.23	.90	.29
15	1.46	1.50	1.50	1.16	.18

$$\mathbf{s}_{\hat{\mathbf{T}}}^{2} = \frac{6}{5} \left[ \frac{1}{6} \sum_{i}^{6} \hat{\mathbf{T}}^{2} - \left( \frac{1}{6} \sum_{i}^{6} \hat{\mathbf{T}} \right)^{2} \right]$$
 (20)

the sum being across the six LOGIST runs. If the T in (20) were normally distributed,  $s_{\hat{T}}^2/\sigma_T^2$  would have an F distribution with 5 and  $\infty$  degrees of freedom.

Since three different items have identical item parameters, the  $\mathbf{s}_{T}^{2}$  for a single item parameter can be averaged across these three items to yield the best available unbiased estimate:

$$\bar{s}_{T}^{2} = \frac{1}{3} \sum_{\Sigma} s_{T}^{2}$$
 (21)

Note that it would be incorrect to pool all 18 values of  $\,T\,$  in an equation like (20), since  $\,\hat{T}\,$  from the same LOGIST run are not independent.

If T  $_{i}$  and S  $_{i}$  represent two different item parameters in the same item

$$\mathbf{\bar{s}}(\hat{\mathbf{T}_i}, \hat{\mathbf{S}_i}) \equiv \frac{1}{3} \stackrel{3}{\times} \mathbf{s}(\hat{\mathbf{T}_i}, \hat{\mathbf{S}_i}) \tag{22}$$

which is the same as (21) except that covariances are substituted for variances. If  $\hat{T}_i$  and  $\hat{S}_j$  represent item parameters in different items, then there are nine different sample covariances to be summed:

$$\bar{\mathbf{s}}(\bar{\mathbf{T}}_{\mathbf{i}}, \bar{\mathbf{S}}_{\mathbf{j}}) = \frac{1}{9} \sum_{i=1}^{33} \mathbf{s}(\hat{\mathbf{T}}_{\mathbf{i}}, \hat{\mathbf{S}}_{\mathbf{j}}) \qquad (23)$$

If T is an ability parameter, (20) still holds. For our purposes, replacing T by  $\Theta$  , we can write

$$\bar{\mathbf{s}}_{\hat{\Theta}}^{2} = \frac{1}{N_{g}} \sum_{\Sigma}^{g} \mathbf{s}_{\hat{\Theta}}^{2} \tag{24}$$

where the sum is over all examinees in group g. When r is at the midpoint of interval g, this average should be roughly equal to the obtained in Section 4.

If subscripts a and b denote different examinees in group  $\boldsymbol{g}$  ,

$$\bar{\mathbf{s}}(\vartheta_{\mathbf{a}},\vartheta_{\mathbf{b}}) = \frac{2}{N_{\mathbf{g}}(N_{\mathbf{g}} - 1)} \sum_{\mathbf{a} \geq \mathbf{b}} \mathbf{s}(\vartheta_{\mathbf{a}}, \vartheta_{\mathbf{b}})$$
 (25)

where the sum is over all pairs of examinees in group  $\,g\,$  . If  $\,a\,$  and  $\,b\,$  denote examinees in groups  $\,g\,$  and  $\,h\,$  respectively (  $\,g\,\neq\,h\,$  ), then

$$\bar{\mathbf{s}}(\mathcal{O}_{\mathbf{a}},\mathcal{O}_{\mathbf{b}}) = \frac{1}{\frac{1}{N_{\mathbf{g}}N_{\mathbf{h}}}} \frac{N_{\mathbf{g}}}{r} \frac{N_{\mathbf{h}}}{r} \mathbf{s}(\mathcal{O}_{\mathbf{a}},\mathcal{O}_{\mathbf{b}}) \qquad (2a)$$

Finally, if  $T_{\underline{\mathbf{i}}}$  is an item parameter and examinee |a| is in group |g| , then

$$\mathbf{\bar{s}}(\mathbf{T_i}, \mathbf{\hat{o}_a}) = \frac{1}{3N_g} \mathbf{\hat{s}} \mathbf{\hat{T}} \mathbf{\hat{s}} \mathbf{\hat{s}} \mathbf{\hat{T}_i}, \mathbf{\hat{o}_a}) \qquad (27)$$

In computing (24) - (27), examinees are grouped on their true values, not on their estimated values.

A problem arises when an examinee obtains a perfect score or a zero score. In this case his  $\varepsilon$  is infinite and cannot be advantageously used. Instead of making some ad hoc adjustment, the 17 examinees for whom this occurred were simply removed from the group of examinees studied, leaving N = 1483. This has the effect of slightly biasing  $\overline{s}$  for the remaining most extreme  $\varepsilon$  values.

#### 8. Numerical Standard Errors

Since the coparameter of an easy item usually cannot be accurately estimated, LOGIST in ordinary use does not estimate them individually. This would prevent the empirical standard errors of Section 7 from agreeing with the theoretical standard errors of Section 4. Since our main purpose is to show that the method of Section 4 can give useful results, the empirical and theoretical standard errors reported here are all estimated or calculated under the condition that the true values of  $c_1$  are known for i=1,2,3,4,5,12. Atoms 1.7 are easy items, item 12 was included because of its low  $a_1$ . For empirical work, the true covalues were supplied to LOGIST, which held them fixed while estimating all other parameters. For theoretical work, the rows and columns of (7) corresponding to  $c_1$ ,  $c_2$ ,  $c_3$ ,  $c_4$ ,  $c_5$ .

and  $c_{12}$  were simply deleted from the information matrix (7) before inversion.

Table 2 compares the empirical standard errors of Section 7 for B with the theoretical standard errors of Section 4. The last three columns show the squared ratios for the three replications of each item; each of these ratios will have an F distribution with 5 and degrees of freedom provided i) B has a normal sampling distribution, ii) B is unbiased, and iii) the theoretical  $c_{\hat{B}}$  from Section 4 is correct. An F above 2.21 or below .229 is significant at the (two-tailed) 10 percent level. Eleven of the ratios are significant. The rumber of ratios less than 1 is approximately the same as the number of ratios greater than 1.

In the past, the only available standard errors for item parameters assumed that the were known. Such standard errors for  $\hat{B}$ , for known —, are given in the second column of the table. A comparison of second and third columns shows very close agreement except for the three easiest items (1,2,3). For these three items, our new theoretical value is larger and agrees better with the empirical value. This gives support to the new theoretical values. The fact that the empirical values (from Section 7) tend to be larger than the theoretical (from Section 4) could be due to n and N not being large enough for asymptotic results. A second likely explanation is that LOGIST was not really run to complete convergence.

Table 3 makes comparisons for A. Again the standard errors of A with a unknown agree closely with the results when is known. The empirical standard errors, although correlating well with the theoretical, seem to be larger. Eleven of the F ratios are

	ëĝ  <sub>∵</sub>	${}^{\circlearrowleft}\widehat{\mathtt{B}}$	$\hat{\mathbf{s}}_{\hat{B}}$			
Item No.	( known)	(Sect. 4)	(Sect. 7)		$s_{\hat{B}}^2/r_{\hat{B}}^2$	
1*	.110	.156	.183	.23	.56	3.34-
2:	.186	.201	.237	1.76	1.49	.93
3*	.045	.071	.063	1.38	.59	.41
4*	.060	.068	.066	.90	. 76	1.17
5*	.100	.099	.103	.37	.40	2.48
6	.125	.121	.131	.28	.63	2.63~
7	.113	.110	.100	1.24	.65	.58
ز	.084	.083	.088	2.31*	.97	.16
9	.055	.055	.067	.37	2.634	1.47
1.0	.069	.069	.106	3.19+	3.62	.33
11	.100	.097	.122	1.45	2.554	.70
12*	.094	.091	.087	.85	1.27	.66
13	.086	.083	.094	1.01	1.20	1.57
14,	.077	.076	.111	1.19	1.49	3.754
15	.072	.075	.093	.40	2.62*	1.65

Significant at 10 percent level.

<sup>\*</sup>The  $\ensuremath{\text{C}}$  parameter for these items is treated as known.

 $\begin{tabular}{ll} \begin{tabular}{ll} Table & 3 \\ \end{tabular} \begin{tabular}{ll} Theoretical and Empirical Standard Errors for & A \\ \end{tabular}$ 

Item No.	à A	Â	s <sub>À</sub>		$\frac{s_A^2}{A}$	
1*	.088	.105	.141	•95	.91	3.60-
2*	.044	.046	.039	.88	.51	.74
3*	.097	.117	.094	1.39	.32	.22÷
4*	.060	.065	.080	. 89	2.77	.86
5*	.045	.047	.054	.63	2.44	.93
6	.103	.102	.123	1.54	.30	2.51
7	.105	.105	.147	1.30	2.25	2.35+
8	.113	.115	.159	1.29	3.20-	1.29
9	.123	.128	.182	1.89	3.39	.80
10	.184	.193	.160	.71	<b>.</b> 55	.79
11	.115	.120	.132	1.42	1.85	.34
12*	.060	.060	.076	.95	2.94 <sup></sup>	.94
13	.151	.157	.187	2.40	1.08	.79
14	.209	.218	.240	1.32	.91	1.43
15	.222	.233	.182	. 25	,65	.93

<sup>\*</sup>Significant at 10 percent level.

<sup>\*</sup>The C parameter for these items is treated as known.

significant. Similar statements apply to Table 4, which shows the comparisons for  $\hat{\mathbf{C}}$  .

Table 5 compares standard errors for  $^3$ . Let us leave column 3 for later discussion. Columns 4 and 5 show standard errors of corresponding to the  $^4$  value in the first column; column 6, however, is computed from (2) for the group of  $^{10}$  people falling in the class interval with midpoint  $^{12}$ . There is good agreement between empirical and theoretical standard errors except for  $^{12}$  < -1.5 . For low  $^{13}$ , asymptotic results do not appear with the usual  $^{11}$  and  $^{11}$  N.

Table 5 shows close agreement of our standard error from Sections

2-4 with the standard error of  $\hat{j}$  when the item parameters are known. The agreement shown here and in previous tables suggests that (1) is a good approximation to the diagonal of (2) and similarly for item parameters, that (2) agrees well with the empirical standard errors.

A comparison of the third and fifth columns in Table 5 shows what happens to -1 when all C must be estimated from the data: For 0 < -1, 0 is sharply affected; for 0 < -2.5, there is very little effect.

Table 6 contains the squared ratios of the empirical standard errors to the theoretical standard errors for the five a closest to the midpoint of the intervals, and within at least .1 of the midpoint. Two of the groups had only two abilities within this restriction. If similar caveats apply as for the item parameters these ratios will have an F distribution with five and a degrees of freedom. Only eight of the ratios are significant at the two-tailed 10% level, and only 16 are greater than 1.

Item No.*	ÖČ! ;	$\frac{\sigma_{\hat{\mathbf{c}}}}{\hat{\mathbf{c}}}$	s <sub>Ĉ</sub>		$\mathbf{s}_{\hat{\mathbf{C}}}^2/\tilde{\mathbf{c}}$	
6	.056	.058	.063	.39	.44	2.79
7	.049	.050	.038	.40	.35	.95
8	.037	.037	.045	3.08+	.76	.43
9	.024	.025	.039	.80	4.71	1.83
10	.025	.026	.034	2.24+	2.68**	.27
11	.036	.037	.043	.98	2.67	.41
13	.026	.027	.037	.89	1.88	2.90
14	.019	.020	.028	2.98	2.55	.43
15	.015	.015	.016	.64	1.23	1.71

<sup>\*</sup>Significant at 10 percent level.

 $<sup>^{*\</sup>text{C}}_1,\dots,^{\text{C}}_5$  , and  $^{\text{C}}_{12}$  are treated as known.

Table 5 . Theoretical and Empirical Standard Errors for  $\ensuremath{^{\mbox{\tiny $\Omega$}}}$ 

		All C	c <sub>l</sub> to	$^{\circ}$ $^{\circ}$ $^{\circ}$ $^{\circ}$ and $^{\circ}$ $^{\circ}$ $^{\circ}$	treated
		unknown		as known	
÷	N <sub>E</sub>	1 ·	A,B,C	·	<del>-</del> 5
-2.75	10	2.090	.951	.966	*
-2.25	35	1.296	.686	.699	1.134
-1.75	93	.861	.516	.525	.797
-1.25	219	.607	.400	.404	.427
75	332	.456	.341	.342	.332
25	326	.349	.295	.295	.279
.25	227	.278	.262	.263	.274
<b>.</b> 75	136	.261	.260	.261	.286
1.25	77	.303	.289	.290	.349
1.75	25	.422	.384	.387	.412
2.25	3	.628	.575	.580	*
2.75	0	.931	.874	.878	*

<sup>\*</sup>Not computed because of small  $N_{\mbox{\scriptsize g}}$  .

Table 6  $\label{eq:fable_form} \textbf{F} \quad \textbf{Ratios for} \quad \hat{\textbf{O}}$ 

9			$\mathbf{s}_{\hat{0}}^{2}/\sigma_{\hat{0}}^{2}$		
-2.75	3.73+	4.41.			
-2.25	.85	.78	.43	11.34	1.16
-1.75	.57	1.90	1.62	.32	18,95
-1.25	.98	.63	.96	.95	.77
<del></del> 75	.26	. 94	.63	.81	.63
25	.71	1.81	.73	• 04	.48
.25	.18+	.98	.74	.80	.77
.75	.61	.35	1.41	1.21	.64
1.25	2.76+	1.82	.98	1.08	1.84
1.75	.67	.41	1.08	1.45	1.78
2.25	.11+	.36			
2.75*					

<sup>†</sup>Significant at 10 percent level.

<sup>\*</sup>There were no  $\theta$  between 2.65 and 2.85.

Table 7 presents the theoretical standard errors of A , B , and C , obtained by the method of Sections 2-4, when all  $C_i$  must be estimated from the data. It is interesting to compare these values with those in Tables 2-4 where  $C_1,\ldots,C_5$ , and  $C_{12}$  were treated as known. We find that the standard errors of  $B_1$  to  $B_5$  are increased drastically by ignorance of  $C_1$  to  $C_5$ ; all other  $\sigma(\hat{B_i})$  are much increased, except for i = 11, 13, and 14. All  $A_i$  show sharply increased standard errors. For items for which  $C_i$  must be estimated, on the other hand, the standard errors of  $C_i$  are little affected by knowledge or ignorance of  $C_1,\ldots,C_5,C_{12}$ . A likely explanation for this is that errors in estimating the scale unit  $B_1$  affect the standard errors of the  $A_i$  and the  $B_i$ , but not of the  $C_i$ .

We have found in Tables 2-7 some illustrative answers to the question: How do estimation errors on one set of items affect the accuracy of estimated parameters for a different set of items? Such effects could not be quantified until now since the standard error of an item parameter estimate was previously known only for fixed .

It is only through the sampling fluctuations of it that estimation errors for one item can affect parameter estimates for another item.

With 18~ C $_{1}$  treated as known, the Fisher information matrix inverted for this study has  $3 \times 45 - 18 + 1498 = 1615$  rows and columns. The matrix inversion by the method of Section 4 used 1232K bytes of memory on an IBM 3031 and took 32 seconds. The computer program dealt with a 45-item test; it did not take advantage of the fact that the 45 items consisted of 3 replicate sets of 15 items each.

Item No.	<u>°à</u>	<u>°</u> À	<u></u>
1	.52	.23	.60
2	2.54	.13	.72
3	.35	.32	.10
4	.26	.15	.14
5	. 97	.10	.32
6	.19	.18	.07
7	,16	.18	.06
8	.14	.21	.041
9	.12	.26	.026
10	.11	.32	.026
11	.10	.18	.039
12	.18	.14	.07
13	.09	.23	.027
14	.08	.31	.020
15	.10	.33	.015

In order to verify the numerical accuracy of the inversion, the information matrix and the variance-covariance matrix were multiplied. The result was an identity matrix accurate to 10 decimal places. The variance-covariance matrix obtained in double precision agreed with the matrix obtained in quadruple precision to all six decimal places printed.

# 9. Sampling Covariances and Correlations

When item parameters are known,  $\frac{1}{a}$  and  $\frac{1}{b}$  (  $a \neq b$  ) are uncorrelated. When ability parameters are known, estimated item parameters for different items are uncorrelated. When both item and ability parameters are estimated, in general all estimates are correlated. The computer printout of the sampling correlations for the present study consists of 10 correlation matrices. These need only be summarized here.

Table 8 shows the theoretical ( T ) and empirical ( E ) correlations between estimates of two different parameters for the same item. The correlations are generally substantial. For comparison, the theoretical correlations when the abilities are known are included. The empirical correlations are obtained by dividing the estimated sampling covariance by the square roots of the estimated sampling variances. If the empirical correlations here have roughly 15 degrees of freedom, their standard error is roughly  $(1-.2)/\sqrt{15} = .26(1-.2)$ . In view of their standard errors, there is very satisfactory agreement of empirical with theoretical correlations.

Table 9 shows both theoretical and empirical correlations for the  $\hat{B}_i$  ( i = 1,2,...,15 ). The corresponding standard errors are

Two Parameter Estimates for the Same Item

ن ،	田						.53	.79	.58	.81	.74	.83		.67	.68	.56
$\rho_{\hat{\mathbf{A}}}^{\hat{\mathbf{A}}}$	T						.70	.71	.67	.60	.61	.77		.70	.59	.54
	AC 9						.76	.76	.72	.65	.61	.77		· (6	.58	.53
Pê	E						.92	.77	.85	.87	.93	.89		99.	.61	18
	1						06.	88.	.81	.67	.68	.74		.59	.42	.21
	- BC   6						.92	06.	.83	69.	69.	75		. 60	55.	.25
ω,	띠	.87	88.	.65	.76	77.	.53	99.	.26	.50	.68	. 70	79	90.	.35	81
Å.	H	98.	.82	.70	.52	.38	.70	79.	.52	.33	.42	.42	51	.21	.03	25
	AB	.82	.80	.55	.42	.35	.73	.67	.56	.37	.41	07.	55	.22	90.	19
	Item	٦	~1	٣	<b>オ</b>	7	9	7	∞	6	10	11	12	13	14	15

6 TABLE

815	-147	-040	-155 -068	218 -039	126	-085	000	-137	-063	-098	-182 -107	005-	-112 -060	000-	(093)
B14	-064	360	348 -032	343 -018	-193	122	-018 005	081 002	-357	-151	-103 -086	078	0-341 0-057	(111)	006
B13	-031	-247	-298 -008	-443 -005	-051	016	156	-062	332	270-043	-011	)-176 )-069	(083	-341	-112
B12	016	-064	007	192 028	086 013	005	-221 -009	101	-129 -013	-137 -052	041	(087	-176 -069	078 -068	005
811	289 -001	029	268 008	-007	236 -009	-153 004	-050	-015	037	1-193 1-035	(122)	041	-011	-103 -086	-182 -107
B10	$-122 \\ 034$	-105	-279 048	-308 029	-205 -004	107	121	025	198	(106)	-193 -035	-137 -052	270-043	-151 -062	-098 -087
B 9	014	-252 -017	-274 -032	-362 -040	046 -062	-041 -051	098 -036	-068 -007	(055)	198 023	037	-129 -013	332 002	-357	-063 -005
B 8	-004 -088	-005	004	038 -089	-072 -088	076 -053	1-120	(088)	-068	025	-015 001	101	-062	081	-137
B 7	-028 -128	126 -069	056	$\frac{-179}{-130}$	-126 -113	014	(100	-120 -042	098 -036	121	-050	-221 -009	156	-018 005	000
B 6	-158	-036	-091 -131	$\frac{-072}{-130}$	)-228 )-117	(131)	014 -062	076	-041 -051	107 -016	-153	002	016	122	-085
B 5	193 158	-092 078	040 151	120	(103	-228	-126 -113	-072 -088	046	-205	236	086	$\frac{-051}{-003}$	-193 001	126 002
B 4	045 334	286 184	308	(068	120 066	-072 -130	-179	038	-362	-308 029	-007	192 028	-443 -005	343 -018	218 -039
В 3	284 509	541	(063	308	040 151	-091 -131	056 -139	004	-274 -032	-279 048	268 008	007	-298 -008	348 -032	-155 -068
8 2	141	(237)	541 284	286 184	-092 078						029	-064		360 -018	-040
8	(183)	141264	284 509	334	193	-158 -124	-028 -128	-00 <b>4</b> -038	014-040	-122 034	28 <b>9</b> -001	016	-031	-064	-147 -050
	ш-	w⊢	<b>ш</b> ⊱-	ш⊢	ш⊢	ш⊢	יש	шH	w⊢	ш⊢	шH	ш⊢	ш⊢	шH	ш⊢
	8 1	B 2	<b>60</b>	7 80	83 22	9 8	8 7	<b>80</b>	89	B 1 0	811	B12	B 1 3	B 1 4	B15
	1 B 2 B 3 B 4 B 5 B 6 B 7 B 8 B 9 B10 B11 B12 B13 B14 B1	B 1 B 2 B 3 B 4 B 5 B 6 B 7 B 8 B 9 B10 B11 B12 B13 B14 B1 1 E (183) 141 284 045 193 -158 -028 -004 014 -122 289 016 -031 -064 -14 7 (156) 264 509 334 158 -124 -128 -088 -040 034 -001 039 -005 -022 -05	B 1 B 2 B 3 B 4 B 5 B 6 B 7 B 8 B 9 B10 B11 B12 B13 B14 B1  1 E (183) 141 284 045 193 -158 -028 -004 014 -122 289 016 -031 -064 -14  7 (156) 264 509 334 158 -124 -128 -088 -040 034 -001 039 -005 -022 -05  2 E 141 (237) 541 286 -092 -036 126 -005 -252 -105 029 -064 -247 360 -04  7 264 (201) 284 184 078 -066 -069 -046 -017 026 007 022 -002 -018 -03	B 1 B 2 B 3 B 4 B 5 B 6 B 7 B 8 B 9 B 10 B 11 B 12 B 13 B 14 B 1 B 1 B 1 B 1 B 1 B 1 B 1 B 1 B	1 E (183) 141 284 045 193 -158 -028 -004 014 -122 289 016 -031 -064 -14 2 E 141 (237) 541 284 045 -036 -036 -046 -017 026 007 022 -005 -022 -05 3 E 284 541 (063) 308 040 -091 056 004 -274 -279 268 007 -298 348 -15 509 284 (071) 377 151 -131 -139 -093 -060 029 003 028 -043 343 21 5 E 045 286 308 (066) 120 -072 -179 038 -362 -308 -007 029 -044 3 343 -21	F (183) 141 284 045 193 -158 -028 -004 014 -122 289 016 -031 -064 -14   F (156) 264 509 334 158 -124 -128 -088 -040 034 -001 039 -005 -022 -05   F (156) 264 509 334 158 -124 -128 -088 -040 034 -001 039 -005 -252 -05   F (156) 264 (201) 284 078 -086 -069 -046 -017 026 007 -279 268 007 -279 360 -034   F 264 (201) 284 184 078 -066 -069 -046 -017 026 007 -279 268 007 -279 348 -15   F 264 (201) 377 151 -131 -139 -093 -040 029 003 028 -443 343 -05   F 334 184 377 (068) 066 -130 -130 -130 -089 -040 029 003 028 -005 -018 -033 -035 046 -019 003 028 -005 -018 -033 001 001 001 001 001 001 001 001 001	F (183) 141 284 045 193 -158 -028 -084 044 014 -122 289 016 -031 -064 -14	F (183) 141   284	F (183) 141 284 045 193 -158 -028 -040 040 034 -081 015 -015 -015 -015 -015 -015 -015 -015	F (183) 141   284   945   193   -158   -028   -046   014   -122   -289   016   -031   -064   -14   -15   -154   -237   -244   509   334   158   -154   -128   -088   -046   015   -025   -015   -052   -055	F (183) 141 284 945 193 -126 -028 -046 034 -061 039 -065 -052 -055 -055 -055 -055 -055 -055 -05	F (184) 141 284 945 193 -154 -084 -044 -045 -287 -289 015 -031 -065 -055 -055 -055 -055 -055 -055 -055	F (183)   141   284   344   193   158   -028   -004   -015   -289   016   -015   -025   -026   -015   -025   -026   -015   -025   -026   -015   -02	F (183) 144   284   945   193   158   -028   -004   -015   -268   015   -015	F (1963) 1441 284 945 193 1156 -1028 -0046 -0146 -135 -289 015 -0151 -0152 -0152 -0153 -0154 -1154 -1154 -1154 -1154 -1155 -0158 -0154 -0155 -

given in parentheses in the diagonal. The only theoretical correlations above .20 are among  $\hat{B}_1$ ,  $\hat{B}_2$ ,  $\hat{B}_3$ , and  $\hat{B}_4$ . These are the four easiest items. Any error in estimating the scale unit  $\bar{B}_1 - \bar{B}_0$  would seriously affect all these items in the same way. It is hard to draw other useful generalizations from this table.

The corresponding table for the  $A_i$  (  $i=1,2,\ldots,15$  ) shows only 3 theoretical correlations above .20:  $\nu_{13}=.27$ ,  $\nu_{14}=.20$ ,  $\nu_{34}=.23$ . With two exceptions (  $\nu_{67}=-.013$ ,  $\nu_{6,12}=-.002$ ), all theoretical correlations are positive.

The highest theoretical correlation among the  $C_i$  ( i = 6,7,..., 11 and 13, 14, 15 ) is  $\rho_{67}$  = .04 . All correlations are positive.

The theoretical correlations between  $A_i$  and  $B_j$  (  $i \neq j$  ) are all below .20 in absolute value, except for items 1-4, which vary from .14 to .38. For  $B_i$  and  $C_j$  (  $i \neq j$ ;  $j \neq 1,2,\ldots,5,12$  ) there are no correlations above .25 in absolute value. For  $A_i$  and  $C_j$ , there are no correlations above .20 in absolute value.

The theoretical correlations between  $\frac{1}{a}$  and  $\frac{1}{b}$  (  $a \neq b$  ) are all less than .04 in absolute value. Between  $\frac{1}{a}$  and  $\frac{1}{b}$ , the largest correlation in absolute value is .15 (when i = 1 and  $\frac{1}{a} = -2.25$ ). Between  $\frac{1}{a}$  and  $\frac{1}{a}$ , the largest is .12 (when i = 1 and  $\frac{1}{a} = -2.25$ ). Between  $\frac{1}{a}$  and  $\frac{1}{a}$ , the largest is .06.

### Summary

When both abilities and item parameters are unknown, the asymptotic sampling variance-covariance matrix developed in this paper appears to provide useful values for the standard errors needed for further research in item response theory. The magnitude of the numerical values in the matrix were very much affected by the method used to define the scale. For a set of artificial data, this variance-covariance matrix compared satisfactorially with empirical results; also with the variance-covariance matrices found by the usual formulas for the case where the abilities are known or where the item parameters are known.

With this matrix, the effect on other items of including items with poorly determined parameters can be studied. Including items with poorly determined c's increases the standard errors of all of the a's and b's but not of the other c's. The effect of different distributions of abilities on the accuracy of item parameters can also be studied. Hopefully a goodness-of-fit test can now be developed for the three-parameter model.

The standard errors of item parameters can now be studied for a situation of common occurrence in equating and item banking: Each of two tests containing common items is administered to a different group of examinees; all parameters are estimated in the same LOGIST run. It is of particular interest to determine how the number of common items affects the standard error of the parameter estimates.

# References

- Bradley, R. A. & Gart, J. J. The asymptotic properties of ML estimators when sampling from associated populations. <u>Biometrika</u>, 1962, <u>49</u>, 205-214.
- Haberman, S. J. Maximum likelihood estimates in exponential response models. The Annals of Statistics, 1977, 5, 815-841.
- Kendall, M. G. & Stuart, A. The advanced theory of statistics (Vol. 1, 3rd ed.). New York: Hafner, 1969.
- Lord, F. M. Applications of item response theory to practical testing problems. Hillsdale, N.J.: Lawrence Erlbaum Associates, 1980.

### DISTRIBUTION LIST

#### Navy

- Provost and Academic Dean
  U.S. Naval Postgraduate School
  Monterey, CA 93940
- Chief of Naval Education and Training Liason Office Air Force Human Resource Laboratory Flying Training Division Williams Air Force Base, AZ 85224
- 1 CDR Mike Curran Office of Naval Research 800 North Quincy Street Code 270 Arlington, VA 22217
- 1 Dr. Pat Federico
  Navy Personnel R & D Center
  San Diego, CA 92152
- Mr. Paul Foley Navy Personnel R & D Center San Diego, CA 92152
- 1 Dr. John Ford
  Navy Personnel R & D Center
  San Diego, CA 92152
- Patrick R. Harrison
  Psychology Course Director
  Leadership and Law Department (7b)
  Division of Professional Development
  U.S. Naval Academy
  Annapolis, MD 21402

- Dr. Norman J. Kerr Chief of Naval Technical Training Naval Air Station Memphis (75) Millington, TN 38054
- 1 Dr. William L. Maloy
  Principal Civilian Advisor for
   Education and Training
  Naval Training Command, Code 00A
  Pensacola, FL 32508
- CAPT Richard L. Martin, USN Prospective Commanding Officer USS Carl Vinson (CNV-70) Newport News Shipbuilding and Drydock Co. Newport News, VA 23607
- 1 Dr. James McBride Navy Personnel R & D Center San Diego, CA 92152
- 1 Mr. William Nordbrock
   Instructional Program Development
   Building 90
   NET-PDCD
   Great Lakes NTC, IL 60088
- 1 Library, Code P201L Navy Personnel R & D Center San Diego, CA 92152

- Commanding Officer
   Naval Research Laboratory
   Code 2627
   Washington, DC 20390
- 1 Psychologist
   ONR Branch Office
   Building 114, Section D
   666 Summer Street
   Boston, MA 02210
- 1 Office of Naval Research
  Code 437
  800 North Quincy Street
  Arlington, VA 22217
- 5 Personnel and Training Research Programs Code 458 Office of Naval Research Arlington, VA 22217
- 1 Psychologist ONR Branch Office 1030 East Green Street Pasadena, CA 91101
- 1 Office of the Chief of Naval Operations Research Development and Studies Branch OP-115 Washington, DC 20350
- The Principal Deputy Assistant Secretary of the Navy (MRA&L) 4E780, The Pentagon Washington, DC 22203
- Director, Research and Analysis Division Plans and Policy Department Navy Recruiting Command 4015 Wilson Boulevard Arlington, VA 22203

- 1 Mr. Arnold Rubenstein Office of Naval Technology 800 N. Quincy Street Arlington, VA 22217
- 1 Dr. Worth Scanland, Director
  Research, Development, Test
   and Evaluation
  N-5
  Naval Education and Training Command
  NAS
  Pensacola, FL 32508
- 1 Dr. Robert G. Smith
   Office of Chief of Naval Operations
   OP-987H
   Washington, DC 20350
- 1 Dr. Alfred F. Smode
   Training Analysis and Evaluation Group
   Department of the Navy
   Orlando, FL 32813
- Dr. Richard Sorensen
  Navy Personnel R & D Center
  San Diego, CA 92152
- 1 Mr. J. B. Sympson Naval Personnel R & D Center San Diego, CA 92152
- l Dr. Ronald Weitzman Code 54 WZ Department of Administrative Services U.S. Naval Postgraduate School Monterey, CA 93940

- 1 Dr. Robert Wisher
  Code 309
  Navy Personnel R & D Center
  San Diego, CA 92152
- l Dr. Martin F. Wiskoff
  Navy Personnel R & D Center
  San Diego, CA 92152
- 1 Mr. John H. Wolfe
  Code P310
  U.S. Navy Personnel Research
   and Development Center
  San Diego, CA 92152
- 1 Mr. Ted M. I. Yellen
   Technical Information Office
   Code 201
   Navy Personnel R & D Center
   San Diego, CA 92152

Army

- 1 Technical Director U.S. Army Research Institute for the Behavioral and Social Sciences 5001 Eisenhower Avenue Alexandria, VA 22333
- 1 Dr. Myron Fisch1
  U.S. Army Research Institute for the
   Social and Behavioral Sciences
  5001 Eisenhower Avenue
  Alexandria, VA 22333
- 1 Dr. Michael Kaplan
  U.S. Army Research Institute
  5001 Eisenhower Avenue
  Alexandria, VA 22333

- 1 Dr. Milton S. Katz
   Training Technical Area
  U.S. Army Research Institute
  5001 Eisenhower Avenue
  Alexandria, VA 22333
- Dr. Harold F. O'Neil, Jr.
  Attn: PERI-OK
  Army Research Institute
  5001 Eisenhower Avenue
  Alexandria, VA 22333
- 1 LTC Michael Plummer
  Chief, Leadership and Organizational
   Effectiveness Division
  Office of the Deputy Chief of Staff
   for Personnel
  Department of the Army
  The Pentagon
  Washington, DC 20301
- 1 Dr. James L. Raney U.S. Army Research Institute 5001 Eisenhower Avenue Alexandria, VA 22333
- 1 Mr. Robert Ross
  U.S. Army Research Institute for
   the Social and Behavioral Sciences
  5001 Eisenhower Avenue
  Alexandria, VA 22333
- 1 Dr. Robert Sasmor
  U.S. Army Research Institute for
   the Social and Behavioral Sciences
  5001 Eisenhower Avenue
  Alexandria, VA 22333
- 1 Commandant
  U.S. Army Institute of Administration
  Attn: Dr. Sherrill
  Ft. Benjamin Harrison, IN 46256
- Dr. Joseph Ward
  U.S. Army Research Institute
  5001 Eisenhower Avenue
  Alexandria, VA 22333

## Air Force

- Air Force Human Resources Laboratory AFHRL/MPD Brooks Air Force Base, TX 78235
- 1 U.S. Air Force Office of Scientific Research Life Sciences Directorate Bolling Air Force Base Washington, DC 20332
- 1 Dr. Earl A. Alluisi
  HQ, AFHRL (AFSC)
  Brooks Air Force Base, TX 78235
- Dr. Genevieve Haddad Program Manager Life Sciences Directorate AFOSR Bolling Air Force Base Washington, DC 20332
- 1 Dr. David R. Hunter
  AFHRL/MOAM
  Brooks Air Force Base, TX 78235
- Research and Measurement Division Research Branch, AFMPC/MPCYPR Randolph Air Force Base, TX 78148
- 1 Dr. Malcolm Ree
  AFHRL/MP
  Brooks Air Force Base, TX 78235

Marines

l Dr. H. William Greenup Education Advisor (E031) Education Center, MCDEC Quantico, VA 22134

- 1 Director, Office of Manpower
   Utilization
  HQ, Marine Corps (MPU)
  BCB, Building 2009
  Quantico, VA 22134
- Special Assistant for Marine
   Corps Matters
  Code 100M
  Office of Naval Research
  800 N. Quincy Street
  Arlington, VA 22217
- MAJ Michael L. Patrow, USMC Headquarters, Marine Corps Code MPI-20 Washington, DC 20380
- 1 Dr. A. L. Slafkosky
   Scientific Advisor
   Code RD-1
   HQ, U.S. Marine Corps
   Washington, DC 20380

Coast Guard

- I Chief, Psychological Research Branch U.S. Coast Guard (G-P-1/2/TP42) Washington, DC 20593
- 1 Mr. Thomas A. Warm
  U.S. Coast Guard Institute
  P.O. Substation 18
  Oklahoma City, OK 73169

Other DoD

1 DARPA
1400 Wilson Boulevard
Arlington, VA 22209

- 12 Defense Technical Information Center
   Cameron Station, Building 5
   Attn: TC
   Alexandria, VA 22314
- Dr. William Graham
  Testing Directorate
  MEPCOM/MEPCT-P
  Ft. Sheridan, IL 60037
- Director, Research and Data
  OASD (MRA&L)
  3B919, The Pentagon
  Washington, DC 20301
- Military Assistant for Training and Personnel Technology Office of the Under Secretary of Defense for Research and Engineering Room 3D129, The Pentagon Washington, DC 20301
- 1 Dr. Wayne Sellman
  Office of the Assistant Secretary
   of Defense (MRA&L)
  2B269 The Pentagon
  Washington, DC 20301

Civil Government

- Mr. Richard McKillip Personnel R & D Center Office of Personnel Management 1900 E Street, NW Washington, DC 20415
- 1 Dr. Andrew R. Molnar Science Education Development and Research National Science Foundation Washington, DC 20550

- Dr. H. Wallace Sinaiko
  Program Director
  Manpower Research and
   Advisory Services
  Smithsonian Institution
  801 North Pitt Street
  Alexandria, VA 22314
- 1 Dr. Vern W. Urry
  Personnel R & D Center
  Office of Personnel Management
  1900 E Street, NW
  Washington, DC 20415
- Dr. Joseph L. Young, Director Memory and Cognitive Processes National Science Foundation Washington, DC 20550

Non-Government

- Dr. James Algina
  University of Florida
  Gainesville, FL 32611
- 1 Dr. Erling B. Andersen Department of Statistics Studiestraede 6 1455 Copenhagen DENMARK
- Psychological Research Unit
  Department of Defense (Army Office)
  Campbell Park Offices
  Canberra, ACT 2600
  AUSTRALIA
- 1 Dr. Isaac Bejar Educational Testing Service Princeton, NJ 08541

- 1 CAPT J. Jean Belanger Training Development Division Canadian Forces Training System CFTSHQ, CFB Trenton Astra, Ontario KOK 1BO CANADA
- 1 Dr. Menucha Birenbaum School of Education Tel Aviv University Tel Aviv, Ramat Aviv 69978 ISRAEL
- I Dr. Werner Birke
   DezWPs im Streitkraefteamt
   Postfach 20 50 3
   D-5300 Bonn 2
   WEST GERMANY
- Dr. R. Darrell Bock
  Department of Education
  University of Chicago
  Chicago, IL 60637
- l Liaison Scientists
  Office of Naval Research
  Branch Office, London
  Box 39
  FPO, NY 09510
- 1 Dr. Robert Brennan
  American College Testing Programs
  P.O. Box 168
  Iowa City, IA 52240
- 1 Dr. C. Victor Bunderson
  WICAT Inc.
  University Plaza, Suite 10
  1160 S. State Street
  Orem, UT 84057
- I Dr. John B. Carroll
  Psychometric Laboratory
  University of North Carolina
  Davie Hall 013A
  Chapel Hill, NC 27514

- 1 Charles Myers Library Livingstone House Livingstone Road Stratford London El5 2LJ ENGLAND
- 1 Dr. Kenneth E. Clark
  College of Arts and Sciences
  University of Rochester
  River Compus Station
  Rochester, NY 14627
- Dr. Norman Cliff
  Department of Psychology
  University of Southern California
  University Park
  Los Angeles, CA 90007
- 1 Dr. William E. Coffman
  Director, Iowa Testing Programs
  334 Lindquist Center
  University of Iowa
  Iowa City, IA 52242
- 1 Dr. Meredith P. Crawford
   American Psychological Association
  1200 17th Street, N
  Washington, DC 20036
- I Dr. Fritz Drasgow
  Yale School of Organization and
   Management
  Yale University
  Box 1A
  New Haven, CT 06520
- I Dr. Mike Durmeyer
  Instructional Program Development
  Building 90
  NET-PDCD
  Great Lakes NTC, IL 60088
- 1 ERIC Facility-Acquisitions 4833 Rugby Avenue Bethesda, MD 20014

- 1 Dr. A. J. Eschenbrenner
  Dept. E422, Bldg. 81
  McDonald Douglas Astronautics Co.
  P.O. Box 516
  St. Louis, MO 63166
- Dr. Benjamin A. Fairbank, Jr.
  McFann-Gray and Associates, Inc.
  5825 Callaghan
  Suite 225
  San Antonio, TX 78228
- l Dr. Leonard Feldt
  Lindquist Center for Measurement
  University of Iowa
  Iowa City, IA 52242
- 1 Dr. Richard L. Ferguson
  The American College Testing Program
  P.O. Box 168
  Iowa City, IA 52240
- Dr. Victor Fields
   Department of Psychology
   Montgomery College
   Rockville, MD 20850
- 1 Univ. Prof. Dr. Gerhard Fischer
  Psychologisches Institut der
   Universitat Wien
  Liebiggasse 5/3
  A 1010 Wien
  AUSTRIA
- l Prof. Donald Fitzgerald University of New England Armidale, New South Wales 2351 AUSTRALIA
- 1 Dr. Edwin A. Fleishman
  Advanced Research Resources Organization
  Suite 900
  4330 East West Highway
  Washington, DC 20014

- 1 Dr. John R. Frederiksen
  Bolt, Beranek, and Newman
  50 Moulton Street
  Cambridge, MA 02138
- Dr. Robert Glaser
  LRDC
  University of Pittsburgh
  3939 O'Hara Street
  Pittsburgh, PA 15213
- 1 Dr. Daniel Gopher
   Industrial and Management Engineering
   Technion-Israel Institute of
   Technology
   Haifa
   ISRAEL
- Dr. Bert Green
  Department of Psychology
  Johns Hopkins University
  Charles and 34th Streets
  Baltimore, MD 21218
- 1 Dr. Ron Hambleton
  School of Education
  University of Massachusetts
  Amherst, MA 01002
- 1 Dr. Delwyn Harnisch University of Illinois 242b Education Urbana, IL 01801
- 1 Dr. Chester Harris School of Education University of California Santa Barbara, CA 93106
- 1 Dr. Lloyd Humphreys
   Department of Psychology
   University of Illinois
   Champaign, IL 61820

- l Library
  HumRRO/Western Division
  27857 Berwick Drive
  Carmel, CA 93921
- Dr. Steven Hunka Department of Education University of Alberta Edmonton, Alberta CANADA
- 1 Dr. Jack Hunter 2122 Coolidge Street Lansing, MI 48906
- 1 Dr. Huynh Huynh
  College of Education
  University of South Carolina
  Columbia, SC 29208
- Prof. John A. Keats
  Department of Psychology
  University of Newcastle
  Newcastle, New South Wales 2308
  AUSTRALIA
- l Mr. Jeff Kelety
  Department of Instructional Technology
  University of Southern California
  Los Angeles, CA 90007
- Dr. Michael Levine
  Department of Educational Psychology
  210 Education Building
  University of Illinois
  Champaign, IL 61801
- 1 Dr. Charles Lewis Faculteit Sociale Wetenschappen Rijksuniversiteit Groningen Oude Boteringestraat 23 9712GC Groningen NETHERLANDS


- l Dr. Robert Linn
  College of Education
  University of Illinois
  Urbana, IL 61801
- Dr. James Lumsden
  Department of Psychology
  University of Western Australia
  Nedlands, Western Australia 6009
  AUSTRALIA
- l Dr. Gary Marco Educational Testing Service Princeton, NJ 08541
- Dr. Scott Maxwell
  Department of Psychology
  University of Houston
  Houston, TX 77004
- 1 Dr. Samuel T. Mayo
  Loyola University of Chicago
  820 North Michigan Avenue
  Chicago, IL 60611
- Prof. Jason Millman
  Department of Education
  Stone Hall
  Cornell University
  Ithaca, NY 14853
- 1 Dr. Melvin R. Novick
  356 Lindquist Center for Measurement
  University of Iowa
  Iowa City, IA 52242
- 1 Dr. Jesse Orlansky
  Institute for Defense Analyses
  400 Army Navy Drive
  Arlington, VA 22202
- 1 Dr. Wavne M. Patience
  American Council on Education
  GED Testing Service, Suite 20
  One Dupont Circle, NW
  Washington, DC 20036

- Portland State University
  P.O. Box 751
  Portland, OR 97207
- 1 Mr. Luigi Petrullo
  2431 North Edgewood Street
  Arlington, VA 22207
- 1 Dr. Diane M. Ramsey-Klee
  R-K Research and System Design
  3947 Ridgemont Drive
  Malibu, CA 90265
- 1 Mr. Minrat M. L. Rauch
  P II 4
  Bundesministerium der Verteidigung
  Postfach 1328
  D-53 Bonn 1
  GERMANY
- Dr. Mark D. Reckase
  Educational Psychology Department
  University of Missouri-Columbia
  4 Hill Hall
  Columbia, MO 65211
- 1 Dr. Andrew Rose
  American Institutes for Research
  1055 Thomas Jefferson St., NW
  Washington, DC 20007
- Dr. Leonard L. Rosenbaum, Chairman Department of Psychology Montgomery College Rockville, MD 20850
- 1 Dr. Ernst 2. Rothkopf Bell Laboratories 600 Mountain Avenue Murray Hill, NJ 07974
- 1 Dr. Lawrence Rudner 403 Elm Avenue Takoma Park, MD 20012

- 1 Dr. J. Ryan
   Department of Education
   University of South Carolina
   Columbia, SC 29208
- Prof. Fumiko Samejima
  Department of Psychology
  University of Tennessee
  Knoxville, TN 37916
- Dr. Kazuo Shigemasu
  University of Tohoku
  Department of Educational Psychology
  Kawauchi, Sendai 980
  JAPAN
- 1 Dr. Edwin Shirkey
   Department of Psychology
   University of Central Florida
   Orlando, FL 32816
- 1 Dr. Robert Smith
   Department of Computer Science
  Rutgers University
  New Brunswick, NJ 08903
- 1 Dr. Richard Snow School of Education Stanford University Stanford, CA 94305
- Dr. Robert Sternberg
  Department of Psychology
  Yale University
  Box 11A, Yale Station
  New Haven, CT 06520
- 1 Dr. Patrick Suppes
  Institute for Mathematical Studies in
   the Social Sciences
  Stanford University
  Stanford, CA 94305

- 1 Dr. Hariharan Swaminathan
  Laboratory of Psychometric and
  Evaluation Research
  School of Education
  University of Massacuusetts
  Amherst, MA 01003
- 1 Dr. Kikumi Tatsuoka
   Computer Based Education Research
   Laboratory
  252 Engineering Research Laboratory
  University of Illinois
  Urbana, IL 61801
- 1 Dr. David Thissen
  Department of Psychology
  University of Kansas
  Lawrence, KS 66044
- 1 Dr. Robert Tsutakawa Department of Statistics University of Missouri Columbia, MO 65201
- 1 Dr. Howard Wainer Educational Testing Service Princeton, NJ 08541
- 1 Dr. David J. Weiss No60 Elliott Hall University of Minnesota 75 East River Road Minneapolis, MN 55455
- l Dr. Susan E. Whitely Psychology Department University of Kansas Lawrence, KS 66044
- 1 Dr. Wolfgang Wildgrube Streitkraefteamt Box 20 50 03 D-5300 Bonn 2 WEST GERMANY

# DATE FILMED



DTIC